

# ***Use of Blood Lactate Measurements in the Critical Care Setting***

**John G Toffaletti, PhD**

**Director of Blood Gas and Clinical Pediatric Labs**

**Professor of Pathology**

**Duke University Medical Center**

**Chief, VAMC Clinical Chemistry Lab**


**Durham, NC**

***email: [john.toffaletti@duke.edu](mailto:john.toffaletti@duke.edu)***


# ***Topics to Discuss***

- **The biochemical mechanisms and clinical processes that can increase blood lactate.**
- **The clinical implications of an increased blood lactate in surgery, ECMO, in the ED, and in sepsis.**
- **The general timing sequence of lactate measurements for monitoring patients in critical care.**
- **The stability of lactate in blood with and without stabilizers.**
- **When and where POC measurements of blood lactate are useful.**

# ***Lactate Testing at Duke Medical Center***



Lactate




# ***Production of Lactate from Pyruvate:***

*Directly Depends on Ratio of NADH/NAD<sup>+</sup>*

*Indirectly Depends on Supply of Oxygen*

Blood



# *The Production of Lactate from Pyruvate Actually Consumes Acid*

<u>Reaction</u>	<u>Net gain/loss of acid</u>
glucose $\longrightarrow$ 2 pyruvate + 2H <sup>+</sup>	produces 2 H <sup>+</sup>
<b>2 pyruvate + 2H<sup>+</sup> <math>\longrightarrow</math> 2 lactate</b>	<b>consumes 2 H<sup>+</sup></b>
ATP + H <sub>2</sub> O $\longrightarrow$ ADP + HPO <sub>4</sub> <sup>=</sup> + H <sup>+</sup>	produces 1 H <sup>+</sup>

See: "Biochemistry of Exercise-Induced Metabolic Acidosis". Am J Physiol Integr Comp Physiol 2004; 287: R502-R516

# ***What Processes Can Elevate Blood Lactate?***

- Normal RBC and muscle cell metabolism: exercise.
- Inadequate oxygen delivered to tissues. Sepsis
- Increased rate of glycolysis: fever. Sepsis
- Decreased rate of clearance or removal:
  - Liver, kidney damage. Sepsis
- Mitochondrial damage from infections and inflammation: Sepsis
  - O<sub>2</sub> radicals, TNF, cytokines, drugs, etc may be involved.

# ***Clinical Uses for Blood Lactate Measurements: Old and New***

- **Monitoring during / after surgery:**
  - open-heart surgery in neonates
  - **adult cardiac operations with CP bypass**
- **Monitoring during ECMO.**
- **Triage use in Emergency Medicine:**
  - **trauma patients, chest pain patients**
  - **criteria for ICU admission.**
- **Detecting / monitoring metabolic alterations in sepsis, septic shock, etc.**



# ***Interpretation of Blood Lactate Results***

- $\leq 1.5$  mmol/L: Normal adult at rest
- 2.2 - 4.0 mmol/L: Moderately elevated
- $> 4.0 - 5.0$  mmol/L: Seriously elevated?
- *But the direction of change may be most important!*

# ***What Does a Blood Lactate Concentration Tell You Clinically?***

- In many patients (surgery, trauma, with sepsis, respiratory distress, etc) an elevation may indicate a problem:
  - insufficient oxygen to tissues, inflammation, etc.
- In an emergency setting with multiple patients to treat:
  - Which patient is sicker?
    - » Which patients can wait for treatment?
    - » Which patients need immediate care?
    - » Which patients are beyond help?
- Is what you are doing making the patient better or worse?

# ***General Format for Using Blood Lactate Measurements***

- **Measure lactate right away:**
  - Lactate normal: GOOD
  - Lactate slightly elevated: Initiate therapy
  - Lactate markedly elevated: Consider more aggressive therapy
- **Measure lactate every 3-6 hours:**
  - Lactate decreasing: GOOD
  - Lactate staying the same: Increase level of therapy
  - Lactate rising: BAD – Consider most aggressive therapy
- **Evaluate after 24 hours:**
  - Lactate normal or close to normal: GOOD
  - Lactate still clearly elevated: Consider more aggressive therapy

# ***Blood Lactate in Pediatric Cardiac Surgery***

# ***Blood Lactate Following Pediatric Cardiac Surgery***


## **Timing of measurements:**

- **Blood lactates are measured after surgery, then every 4-8 hrs after as necessary during recovery.**

## **Interpretation:**

- **Post-surgery lactate of  $\geq 4$  mmol/L generally indicates more intensive care will be needed.**
- **A definite rise in lactate at any time warrants immediate intervention.**
- **After 24 hours, lactate should be normalizing.**


## ***Pediatric Open-Heart Surgery: Closure of Ductus Arteriosus with Placement of Shunt from Aorta to Pulmonary Artery***



# ***Blood Lactate Use in ECMO***

***(Extracorporeal Membrane Oxygenation)***

# ***ECMO = Extracorporeal Membrane Oxygenation***





# ***Interpretation of Blood Lactate Results During ECMO***

- In questionable cases, lactate measurement can help determine if patient goes on ECMO or not ( $\geq 5$  mmol/L).
- Lactate declining or remaining low during ECMO is good.
- If lactate increases or remains elevated:
  - may increase pump flow, blood volume, or hematocrit.
  - evaluate for cardiac problems.
  - consider changing to veno-arterial ECMO.

***Blood Lactate in Adult  
Cardiopulmonary Bypass  
Surgery***

# ***Information Provided by Blood Lactate Measurements In Adult Cardiopulmonary Bypass (CABG) Surgery***

**Monitoring blood lactate evaluates the complex metabolic state of the patient recovering from cooling, hemodilution, anesthesia, vasoactive drugs, inflammation, coagulopathies, etc.**

# ***Principles of Evaluating an Elevated Lactate After Open-Heart Surgery***

- If reperfusion is good, lactate should decline by 1-2 hours after surgery.
  - However, lactate declines slowly in some patients.
- If lactate remains elevated 1-2 hr after surgery:
  - Make sure cardiac output is good.
  - Make sure airways are clear.
  - Evaluate liver function
    - » liver shutdown can diminish lactate removal.
  - Look for gut ischemia or peripheral ischemia.

# Case 1: CABG Operation with No Complications

## 67 yo male; recent Myocardial Infarction

Time	8:40	9:15	10:00	11:15	11:30	12:00	14:00
FI-O <sub>2</sub>	0.40	0.40	0.70	0.70	0.21 (RA)	0.21	1.00
pO <sub>2</sub>	108	101	210	280	180	45	120
%O <sub>2</sub> Hb	98.5	96.7	99.2	99.6	99.3	84.0	98.8
Hb	11.5	10.8	8.2	8.0	8.2	8.5	10.2
O <sub>2</sub> content	15.7	14.5	11.3	11.1	11.3	9.9	14.0
Lactate	1.2	0.9	1.5	2.5	3.8	4.6	2.5

*Patient on pump*

Rise in lactate  
post-op is a relatively  
normal occurrence.

## ***Case 2: CABG Patient With Post-Operative Complications***


- 56 year old male underwent open-heart surgery for coronary artery bypass.
- Blood lactates were measured:
  - Lactate during surgery was 3.2 mmol/L.
  - 4 hr post-surgery lactate was 6.1 mmol/L.
- Several parameters were re-checked:
  - Cardiac output was good
  - No evidence of gut ischemia
  - No problems with breathing
  - Poor peripheral pulses were noted in leg.

## ***Case: CABG Patient Post-Op (cont'd)***

- Patient had an intra-aortic balloon pump inserted through femoral artery to increase cardiac output post-op.
  - Balloon pump may be constricting blood flow to leg.
- Balloon pump was removed from femoral artery.
- Lactate measured 2 hours later was 1.7 mmol/L (normalizing).



# ***Use of Lactate in ED for Trauma and Hypovolemic Shock***






# ***Early Report on Value of Blood Lactate Measurements in Trauma Patients***

- A study of 76 patients admitted to the ICU from either the OR or the ED found that the **time needed to normalize blood lactate** predicted survival rate of patients:
  - 100% (27 of 27) survival when lactate normalized in 24 hours.
  - 78% (21 of 27) survived when lactate normalized within 24-48 hours.
  - 14% (3 of 22) survived if lactate did not normalize by 48 hours.

Abramson, et al: J Trauma 1993; 35: 584-589.

# ***Lactate As Predictor of Survival in Trauma Patients***



## ***Blood Lactate Is Also Helpful in ED for Treating Hypovolemic Shock***

- For hypovolemic shock from:
  - Bleeding, dehydration, etc.
  - Cardiogenic shock
- If resuscitation attempts decrease lactate:
  - Continue on this course.
- If blood lactate stays the same or increases:
  - Look for other causes: sepsis, etc.

# Sepsis



# ***What Is Sepsis and its Progression to More Severe Stages?***


- **Sepsis is an overwhelming response to a systemic infection:**
  - Has SIRS criteria + infection.
- **Severe Sepsis is when a severe infection causes organs to start failing.**
  - May progress to MODS (multiple organ dysfunction syndrome)
- **Sepsis may progress to Septic Shock:**
  - profound drop in blood pressure,
  - organ dysfunction,
  - frequently death (*but EGDT is beneficial!*)

# ***What Are the Criteria for Systemic Inflammatory Response Syndrome (SIRS)?***

- Heart rate > 90/min
- Respiratory rate > 20/min (or  $p\text{CO}_2 < 32$  mmHg).
- Temperature < 35 or > 38 °C
- WBC > 12,000 or < 4,000/mm<sup>3</sup> or > 10% Bands.

*(These are very non-specific criteria for sepsis)*


# *Timeline of Events as Infection Progresses to Sepsis, Septic Shock, and MODS*



SIRS = Systemic Inflammatory Response Syndrome

MODS = Multiple Organ Dysfunction Syndrome

# EARLY GOAL-DIRECTED THERAPY: RESULTS





# ***3- and 6-Hour Bundles in Goal Directed Therapy Protocol for Sepsis in the ED***

- **3-Hour Bundle to rapidly identify patients likely to have sepsis:**
  - Order arterial or mixed venous **lactate**.
  - Order blood cultures.
  - Administer broad spectrum antibiotics.
  - Give fluid bolus if hypotensive or lactate  $>4$  mmol/L.
  - Order CBC, urinalysis, CAT scans, X-rays, etc as appropriate.
  - **Measuring procalcitonin may have great value here!**
- **6-Hour Bundle:**
  - Administer vasopressors if BP is low and unresponsive to fluids.
  - Adjust antibiotics if blood culture results available.
- **Options if hypotension persists and/or lactate remains  $\geq 4$  mmol/L:**
  - Give red cells to achieve  $s_{cv}O_2 \geq 70\%$  or  $s_vO_2 \geq 65\%$
  - Consider mechanical ventilation.


# ***Lactate Measurements in ED for Evaluating Sepsis***

**An elevated lactate in sepsis suggests several possibilities:**

- **Inadequate O<sub>2</sub> delivery:**
  - » Hypovolemia
  - » Shock
  - » Circulatory abnormality: vasoconstriction/vasodilation
- **Problem with O<sub>2</sub> utilization:**
  - » Mitochondrial dysfunction

# ***There Are Many Ways to Lose Your Mitochondria***

## ***By Drugs, Cytokines, Oxygen Radicals***




## ***Increase in Lactate (mmol/L) in Blood Containing No Additive or Fluoride / Oxalate***

<u>Sample</u>	<u>Temp</u>	<u>Additive</u>	<u>Time</u>	Mean increase <u>in Lactate (mmol/L)</u>
plasma	4-23 °C	F / Ox	8 h	< 0.03
plasma	RT	none	2 h	0.10
WB	RT	F / Ox	2 h	0.10
WB	ice	none	60 min	0.10
WB	RT	none	30 min	0.30 (~1%/min)

From Westgard, Clin Chem 1972; Toffaletti, Clin Chem 1992;  
and Astles, Clin Chem 1994

# ***Lactate Changes in Heparinized Blood Gas Samples at Room Temp***




# ***Production of Lactate from Pyruvate:***

*Directly Depends on Ratio of NADH/NAD<sup>+</sup>*

*Indirectly Depends on Supply of Oxygen*

Blood



# ***Summary of Issues with Blood Lactate Measurements***

- Mechanisms for elevated lactate are being clarified.
- Recommendation of lactate testing in EGDT has markedly increased test usage.
- Lactate is becoming a marker for overall mitochondrial damage.
- When to measure and how to interpret?
  - Well established for peds open-heart and ECMO.
  - Becoming established for sepsis, triage in ED, and adult open-heart surgery.


# ***Potential Areas for POC or Lab Measurements of Lactate***

<b>Location</b>	<b>TA-Time Needed</b>	<b>POC</b>	<b>Near Pt Lab</b>	<b>Central Lab</b>
Emergency Dept.	~30 min	ED very chaotic (?)	YES (\$)	May be acceptable
Open-Heart Surgery	5-15 min	YES	YES (\$)	NO
ECMO	5-30 min	YES (but low test #'s)	YES (\$)	May be acceptable
Sepsis	60 min	YES (but many areas to cover)	YES (\$)	Acceptable

**(\$)** = Other tests and test volumes necessary to justify a near-patient laboratory.



# ***Lactate Testing at Duke Medical Center***



***Thank you for your attendance  
and attention!***